

Alternatives To Opiates (ALTOSM) Program

Alexis LaPietra, DO
Medical Director of Emergency Medicine Pain Management
St. Joseph's Regional Medical Center
Paterson, NJ

ALTOSM CLINICAL APPLICATIONS

Renal Colic

1. Toradol 30 mg IV
2. Cardiac Lidocaine 1.5 mg/kg IV in 100 mL NS over 10 minutes (MAX 200 mg)
3. Acetaminophen 1000 mg PO
4. 1 L 0.9% NS bolus

Musculoskeletal Pain (sprains, strains, or opiate naïve lower back pain)

1. Acetaminophen 1000 gm PO
2. Motrin 600 mg PO OR Toradol 30 mg IV/IM
3. Muscle Relaxant (choose one of the following)
 - a. Flexeril 5 mg PO (patients >65 years old OR <70 kg OR concerns for somnolence)
 - b. Flexeril 10 mg PO (patients >70 kg)
 - c. Valium 5 mg PO
4. Lidoderm patch to most painful area, MAX 3 patches instruct patient to remove after 12 hours
5. Gabapentin (neuropathic component of pain)
 - a. 300 mg PO
6. Trigger Point Injection with 1-2 mL of Marcaine 0.5% or Lidocaine 1%

Acute on Chronic Radicular LBP (opiate tolerant)

1. Acetaminophen 1000 mg PO
2. Motrin 600 mg PO OR Toradol 30 mg IV/IM
3. Muscle Relaxant (choose one of the following)
 - a. Flexeril 5 mg PO (patients >65 years old OR <70 kg OR concerns for somnolence)
 - b. Flexeril 10 mg PO (patients >70 kg)
 - c. Valium 5 mg PO

4. Gabapentin (neuropathic component of pain)
 - a. 300 mg PO
5. Dexamethasone 8 mg IV
6. Lidoderm patch to most painful area, MAX 3 patches instruct patient to remove after 12 hours
7. Trigger Point Injection with Marcaine 0.5% or Lidocaine 1% 1-2 mL
8. Ketamine 0.1-0.3 mg/kg in 50 cc NS over 10 minutes
 - a. Ketamine 0.1 mg/kg/hour until pain is tolerable

Headache

1. Reglan 10 mg PO/IV
2. 1 L 0.9% NS bolus
3. Motrin 600 mg PO Or Toradol 30 mg IM/IV
4. Tylenol 1000 mg PO
5. Cervical or Trapezius Trigger Point Injection with Marcaine 0.5% or Lidocaine 1%
If <50% pain relief then
6. Magnesium 1 gm IV over 60 minutes
7. Valproic Acid 500 mg/50 cc NS over 20 mins
8. Dexamethasone 4-8 mg IV
If <50% pain relief then
9. Haldol 5 mg IV
If <50% pain relief then place in OBS unit with Neuro consult

Extremity Fracture or Joint Dislocation

(Steps 1-3 done while setting up for block)

1. Ketamine Intranasal 0.5 mg/kg (concentration 50 mg/mL)
 - a. MAX dose 50 mg; MAX volume per nare 1 mL
2. Nitrous Oxide titrate up to 70%
3. Tylenol 1000 mg PO
4. Ultrasound Guided Regional Anesthesia
 - a. Joint Dislocation
 - i. Lidocaine 0.5 % peri-neural infiltration (MAX 5 mg/kg)
 - b. Extremity Fracture
 - i. Ropivacaine 0.5% peri-neural infiltration (MAX 3 mg/kg)

References

1. Balakrishnamoorthy R, Horgan I, Perez S, Steele MC, Keijzers GB. Does a single dose of intravenous dexamethasone reduce Symptoms in Emergency department patients with low Back pain and RADiculopathy (SEBRA)? A double-blind randomised controlled trial. *Emerg Med J.* 2015 Jul;32(7):525-30.
2. Blaivas M, Adhikari S, Lander L. A prospective comparison of procedural sedation and ultrasound-guided interscalene nerve block for shoulder reduction in the emergency department. *Acad Emerg Med.* 2011 Sep;18(9):922-7.
3. Colman I, Friedman BW, Brown MD, Innes GD, Grafstein E, Roberts TE, Rowe BH. Parenteral dexamethasone for acute severe migraine headache: meta-analysis of randomised controlled trials for preventing recurrence. *BMJ.* 2008 Jun 14;336(7657):1359-61.
4. [Cleveland Clinic Algorithm \(ctrl and click\)](#)
5. Friedman BW, Dym AA, Davitt M, Holden L, Solorzano C, Esses D, Bijur PE, Gallagher EJ. Naproxen With Cyclobenzaprine, Oxycodone/Acetaminophen, or Placebo for Treating Acute Low Back Pain: A Randomized Clinical Trial. *JAMA.* 2015 Oct 20;314(15):1572-80.
6. Fahmida Ghaderibarmi, Nader Tavakkoli, Mansoureh Togha.. Intravenous Valproate versus Subcutaneous Sumatriptan in Acute Migraine Attack. *Acta Medica Iranica* 2015. 53(10):633-636
7. Ferrini R, Paice JA. .How to initiate and monitor infusional lidocaine for severe and/or neuropathic pain. *J Support Oncol.* 2004 Jan-Feb;2(1):90-4
8. Galer BS, Gammaitoni AR, Oleka N, Jensen MP, Argoff CE. Use of the lidocaine patch 5% in reducing intensity of various pain qualities reported by patients with low-back pain. *Curr Med Res Opin.* 2004;20 Suppl 2:S5-12.
9. Gelfand AA, Goadsby PJ. A Neurologist's Guide to Acute Migraine Therapy in the Emergency Room. *The Neurohospitalist.* 2012;2(2):51-59.
10. Herres J, Chudnofsky CR, Manur R, Damiron K, Deitch K. The use of inhaled nitrous oxide for analgesia in adult ED patients: a pilot study. *Am J Emerg Med.* 2016 Feb;34(2):269-73.
11. Kranke P, Jokinen J, Pace NL, Schnabel A, Hollmann MW, Hahnenkamp K, Eberhart LH, Poepping DM, Weibel S. Continuous intravenous perioperative lidocaine infusion for postoperative pain and recovery. *Cochrane Database Syst Rev.* 2015 Jul 16;7.

12. Linde M, Mulleners WM, Chronicle EP, McCrory DC. Valproate (valproic acid or sodium valproate or a combination of the two) for the prophylaxis of episodic migraine in adults. *Cochrane Database Syst Rev.* 2013 Jun 24;6
13. Moore RA, Derry S, Wiffen PJ, Straube S, Aldington DJ. Overview review: Comparative efficacy of oral ibuprofen and paracetamol (acetaminophen) across acute and chronic pain conditions. *Eur J Pain.* 2015 Oct;19(9):1213-23
14. Cohen V, Motov S, Rockoff B, Smith A, Fromm C, Bosoy D, Hossain R, Likourezos A, Jellinek-Cohen SP, Marshall J. Development of an opioid reduction protocol in an emergency department. *Am J Health Syst Pharm.* 2015 Dec 1;72(23):2080-6.
15. Soleimanpour H, Hassanzadeh K, Vaezi H, Golzari SE, Esfanjani RM, Soleimanpour M. Effectiveness of intravenous lidocaine versus intravenous morphine for patients with renal colic in the emergency department. *BMC Urol.* 2012 May 4;12:13.
16. Singh A, Alter HJ, Zaia B.. Does the addition of dexamethasone to standard therapy for acute migraine headache decrease the incidence of recurrent headache for patients treated in the emergency department? A meta-analysis and systematic review of the literature. *Acad Emerg Med.* 2008 Dec;15(12):1223-33
17. Vigneault L, Turgeon AF, Côté D, Lauzier F, Zarychanski R, Moore L, McIntyre LA, Nicole PC, Fergusson DA. Perioperative intravenous lidocaine infusion for postoperative pain control: a meta-analysis of randomized controlled trials. *Can J Anaesth.* 2011 Jan;58(1):22-37.